

()

*

(// : // :)

()

"

() ×
(P < /)

() (P < /)
(P > /)
(P < /)

(P < /)

(P > /)

(P > /)

T4 T3

()

:

.2005)

(Nabizadeh et al.,

(Newkirk et al., 2003)

/
(Tripathi & Mishra, 2007)

)

(

.(/ /)

OO

(Bell et al., 1999)

(NRC,

1994)

(Tahir et al., 2005)

(Vahjen et al., 2005)

(AOAC, 2002)

(Meng & Slominski, 2005)

(Mushtaq et al., 2007)

... :
" .
: ((T₄ T₃)
/ / REF KT2CT
)
(

.(Mawson et al., 1993)

() ×
.(Smith & Campbell, 1976) ()
()
.(SAS, 1997)

.(Yapar & Clandinin, 1972)

(P< /)
%
.(Zokaeinia, 2004)

()
()
()
(P< /)
()
()

(Summers et al., 1989)

()

/	/	/	/	/	/	/	/	/	/	(%)
		/	/	/	/			/	/	(%)
/	/	/	/	/	/	/	/			(% /)
										(%)
/	/	/	/	/	/					(%)
/	/	/	/	/	/	/	/	/	/	(%)
/	/	/	/	/	/	/	/	/	/	(%)
/	/	/	/	/	/	/	/	/	/	(%)
/	/	/	/	/	/	/	/	/	/	(%)
/	/	/	/	/	/	/	/	/	/	(%)
						/	/			(ppm)
/	/	/	/	/	/	/	/	/	/	(%)

()

/	/	/	/	/	/	/	/	/	/	(%)
		/	/	/	/	/	/	/	/	(%)
/	/	/	/	/	/	/	/			(% /)
										(%)
		/	/	/	/	/	/			(%)
		/	/	/	/	/	/	/	/	(%)
		/	/	/	/	/	/	/	/	(%)
/	/	/	/	/	/	/	/	/	/	(%)
/	/	/	/	/	/	/	/	/	/	(%)
/	/	/	/	/	/	/	/	/	/	(%)
						/	/			(ppm)
/	/	/	/	/	/	/	/	/	/	(%)

%
% + %
% + %
% + %
%

%
% + %
% + %
% + %
%

(Tripathi & Mishra, 2007)

(Zokaenia, 2004)

	T ₄		T ₃		
()	()	()	()	()	()
/ a	/	/	/	/	/
/ b	/	/	/	/	/
/ b	/	/	/	/	/
/ c	/	/	/	/	/
/ c	/	/	/	/	/
/	/	/	/	/	SEM
/ a	/ b	/	/	/	/
/ a	/ a	/	/	/	/
/	/	/	/	/	SEM

(Pearson et al., 1989; Zokaenia, 2004; Giffiths et al., 1980)

(Olomu et al., 1975)

(Karujewa et al., 1990)

(P < /) .

() ()

%

(%)	(%)	(%)	(%)	(%)	
/ a	/ b	/	/ a	/	/
/ b	/ ab	/	/ a	/	/
/ b	/ ab	/	/ b	/	/
/ b	/ a	/	/ b	/	/
/ ab	/ a	/	/ a	/	/
/	/	/	/	/	SEM
/ a	/ a	/ a	/ a	/	/
/ a	/ a	/ a	/ a	/	/
/	/	/	/	/	SEM

(P < /) .

()

()

REFERENCES

1. Association of Official Analytical Chemists. (2002). Official Methods of Analysis. 16th ed. Association of Official Analytical Chemists, Washington DC. USA.

2. Bell, J. M., Rokow, G. & Downey, R. K. (1999). Mineral composition of oil free seeds Brassica napus, B. rapa, and B. Juncea as affected by location and year, *Canadian Journal of Animal Science*, 79(4), 405-408.
3. Giffiths, N. M., Fenwick, G. R. Pearson, A. W. Greenwood, N. M. & Butler, E. J. (1980). Effects of rapeseed meal on broilers studies of meat flavor, liver haemorrhage and trimethylamine oxidase activity. *Journal of Science of Food and Agriculture*, 31(2), 188-193.
4. Karujewa, H., Ijabuji, E. G. & Reece, R. L. (1990). Effect of dietary level of rapeseed meal and polyethyleneglycole on the performance of male broilers. *Poultry Science*, 31(3), 345-555.
5. Mawson, R., Heany, R. K. Zdunczyk, Z. & Kozłowska, H. (1993). Rapeseed meal glucosinolates and their antinutritional effects on flavour and palatability. *Die Nahrung*, 37: 36-344.
6. Meng, X., & Slominski, B. A. (2005). Nutritive values of corn, soybean meal, canola meal, and peas for broiler chickens as affected by a multicarbohydrase preparation of cell wall degrading enzymes, *Poultry Science*, 84(8), 1242-1251.
7. Mushtaq, M., Sarwar, G., Ahmad, M. A., Mirza, H., Nawaz, M. M., Mushtaq, H. & Noreen, U. (2007). Influence of canola meal-based diets supplemented with exogenous enzyme and digestible lysine on performance, digestibility, carcass, and immunity responses of broiler chickens. *Poultry Science*, 86(10), 2144 - 2151.
8. Nabizadeh, A., Nasiri Moghaddam, H. & Shahroudi, F. E. (2005). Effect of fish meal protein substitution with blood meal on broiler performance. *Iranian Journal of Agricultural Science*, 36(3), 273-280 (In Farsi).
9. Newkirk, R. W., Classen, H. L., Scott, T. A. & Edney, M. J. (2003). The digestibility and content of Amino acid in toasted and non-toasted canola meal. *Canadian Journal of Animal Science*, 38(1), 131-139.
10. National Research Council, 1994. Nutrient Requirements for Poultry, 9th Rev. Ed. NY.
11. Olomu, J. M., Robblee, A. R. & Clandinin, D. R. (1975). Effect of span rapeseed on the performance organ weights and composition of the carcass, heart and liver of broiler chicks, *Poultry Science*, 54(5), 722-726.
12. Pearson, A. W., Greenwood, N. M., Butler, E. J. & Fenwick, G. R. (1983). Biochemical changes in layer and broiler chickens when fed on a high glucosinolates rapeseed meal. *British Poultry Science*, 24(3), 417-427.
13. SAS Institute. 1997. SAS/State users Guide. Version 6.03 SAS Institute Inc. Carry, NC.
14. Smith, T. K. & Campbell, L. D. (1976). Rapeseed meal glucosinolates metabolism and effect on performance of laying hens. *Poultry Science*, 55(7), 861-867.
15. Summers, J. D., Bedford, M. & Spratt, M. (1989). Amino acid supplementation of canola meal. *Canadian Journal of Animal Science*, 69(3), 469-475.
16. Tahir, M., Saleh, F., Ohtsuka, A. & Hayashi. K. (2005). Synergistic effect of cellulase and hemicellulase on nutrient utilization and performance in broilers fed a corn–soybean meal diet. *Animal Science Journal*, 76(6), 559-565.
17. Tripathi, M. K. & Mishra, A. S. (2007). Glucosiolates in animal nutrition- A review. *Animal Feed Science and Technology*, 132(1,2), 1-27.
18. Vahjen, W., Busch, T. & Simon, O. (2005). Study on the use of soyabean polysaccharide degrading enzymes in broiler nutrition. *Animal Feed Science and Technology*, 120(3,4), 259–276.
19. Yapar, Z. & Clandinin, D. R. (1972). Effect of tannis in rapeseed meal on its nutritional value for chicks. *Poultry Science*, 51(2), 222-228.
20. Zokaenia, F. (2004). *Effect of different levels of canola meal with phytase enzyme on broiler performance*. MSc dissertatation. Isfahan University of Technology, Isfahan, Iran (In Farsi).

